

PATIENT INFORMATION
ENBUMYST (En-byoo-mist)
(bumetanide nasal spray)

What is the most important information I should know about ENBUMYST?

- **ENBUMYST is a diuretic** that causes fluid loss and may lead to abnormalities with blood electrolytes and body energy systems (metabolic). Too much fluid loss can lead to dehydration and decreased blood volume and increase your risk of blood clots. Blood abnormalities may include decreased salts (electrolytes), and increased nitrogen, glucose and uric acid. The chance of getting these abnormalities is higher in people who are elderly, use higher doses, or who do not get enough electrolytes by mouth. Your healthcare provider should do blood tests to check for these abnormalities during treatment with ENBUMYST.
- **You should not use ENBUMYST if you have severe electrolyte loss until it gets better.**

Tell your healthcare provider if you get any of the following signs or symptoms:

- unusual thirst
 - dizziness or lightheadedness
 - weakness or tiredness
 - fast heart rate
 - dry mouth or tongue
 - sudden pain, redness or swelling in your joints, especially your big toe (gout)
 - pain, swelling, redness, or warmth in your leg
 - shortness of breath
 - fast breathing
 - high blood sugar
 - dark urine
 - skin is pale, tight, cool or clammy
 - muscle cramps
 - confusion
- See **“What are the possible side effects of ENBUMYST?”** for more information about side effects.

What is ENBUMYST?

ENBUMYST is a prescription medicine used in adults to treat swelling caused by too much fluid in the body (edema) from congestive heart failure, liver disease and kidney disease, including a kidney problem called nephrotic syndrome. It is not known if ENBUMYST is safe and effective in children.

Who should not use ENBUMYST?

Do not use ENBUMYST if you:

- have very low or no urine production
- are allergic to bumetanide or to any of the ingredients in ENBUMYST. See the end of this Patient Information for a complete list of ingredients in ENBUMYST.

Before using ENBUMYST, tell your healthcare provider about all your medical conditions, including if you:

- have low levels of salts (electrolytes) in your blood, or are on a low-salt diet
- have diarrhea
- have diabetes
- have nasal problems including a runny nose, nasal congestion, nasal polyps, history of injury such as a broken nose, or any past nasal surgery
- are pregnant or plan to become pregnant. It is not known if ENBUMYST will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if ENBUMYST passes into your breast milk. If you use ENBUMYST and breastfeed, watch your baby for signs of urinating more than usual, dehydration, or unusual sleepiness, drowsiness, or sluggishness.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

ENBUMYST and other medicines may affect each other, causing side effects, and may affect the way each other works.

Especially tell your healthcare provider if you take or use:

- other nasal sprays
- water pills (diuretics)
- lithium
- probenecid
- indomethacin
- antibiotic medicines called aminoglycosides
- medicines to treat high blood pressure

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I use ENBUMYST?

- **Read the Instructions for Use that come with ENBUMYST** for detailed information about how to use the device.
- Use ENBUMYST exactly as your healthcare provider tells you to. Talk to your healthcare provider or pharmacist if you have questions about how to use ENBUMYST.
- **ENBUMYST is for use in the nose only.**
- Each ENBUMYST device has 1 spray of medicine and cannot be reused. Do not test or prime (pre-spray) the device.
- ENBUMYST comes in a carton with 12 devices. Use the following number of nasal spray devices for your prescribed dose:

Prescribed Dose	Number of Nasal Spray Devices
0.5 mg	1 device
1 mg	2 devices
1.5 mg	3 devices
2 mg	4 devices

- Your healthcare provider will prescribe the dose of ENBUMYST that is right for you.
- ENBUMYST is for short term use only for a single treatment course (episode) of edema.
- If you use too much ENBUMYST call your healthcare provider right away.

What are the possible side effects of ENBUMYST?

ENBUMYST may cause serious side effects, including:

- **See “What is the most important information I should know about ENBUMYST?”**
- **Worsening kidney function.** Your healthcare provider may stop treatment if your kidney function gets worse.
- **Ear damage (ototoxicity)** including ringing in the ears (tinnitus) or hearing loss, especially in people with severe kidney problems, taking higher than recommended doses, with low levels of protein in the blood, and when used with certain other medicines.

The most common side effects of ENBUMYST include:

- low levels of fluid in the body
- headache
- muscle cramps
- dizziness
- low blood pressure
- nausea
- abnormal brain functioning in people with certain liver problems

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of ENBUMYST.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store ENBUMYST?

- Store ENBUMYST at room temperature between 59°F to 77°F (15°C to 25°C).
- **Do not freeze.**

Keep ENBUMYST and all medicines out of the reach of children.

General information about the safe and effective use of ENBUMYST.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ENBUMYST for a condition for which it was not prescribed. Do not give ENBUMYST to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about ENBUMYST that is written for health professionals.

What are the ingredients in ENBUMYST?

Active Ingredient: bumetanide

Inactive Ingredients: benzyl alcohol (0.5 mg per 0.1 mL spray), carboxymethylcellulose sodium, mannitol, potassium hydroxide (pH modifier), hydrochloric acid (to adjust pH) and water for injection.

Manufactured for Corstasis Therapeutics, 3535 Executive Terminal Drive, Henderson, NV 89042

Copyright © 2025 Corstasis Therapeutics. All rights reserved.

For more information about ENBUMYST, call **1-877-300-5339** or visit ENBUMYST.com.